

Graphdiyne: A new member of carbon family

Jin ZHANG

Center for Nanochemistry, College of Chemistry and Molecular Engineering,
Peking University, Beijing 100871, China.

Email: jinzhang@pku.edu.cn

Graphdiyne (GDY) is an ordered two-dimensional (2D) carbon allotrope comprising sp and sp^2 hybridized carbon atoms with high degrees of π -conjugation, which features a natural bandgap and superior electric properties. However, the synthesis of well-defined GDY remains challenging due to the free rotation around alkyne-aryl single bonds and the lack of thickness control. Herein, we developed several rational approaches to synthesize high-quality structure-controlled GDY. We first demonstrated that the morphology of GDY could be finely controlled by using a modified Hay-Glaser coupling reaction under optimized reaction conditions. Unique vertically grown γ -GDY nanowalls (~ 200 nm) were fabricated on either copper foils or foams^[1,2]. β -GDY, another new member of graphyne family, was also explored using similar method with tetraethynylethene monomers^[3]. Notably, we recently reported a facile synthetic route to synthesize ultrathin single-crystalline GDY, through a solution phase van der Waals (vdW) epitaxial strategy^[4]. The as-grown GDY film has a tri-layer structure with a ABC stacking order as directly observed by electron microscopy. The high quality of the as-grown GDY film and the graphene-enhanced Raman scattering (GERS) effect ensure the predicted Raman fingerprints belonging to a perfectly ordered 2D GDY structure^[5] are experimentally observed. Finally, encouraged by the intriguing properties of such 2D acetylenic carbon allotropes, we designed various GDY-based hierarchical architectures and composites towards practical applications. As one example, a three-dimensional (3D) GDY foam was synthesized and further used for oil/water separation, exhibiting both high efficiency and good recyclability^[2]. Considering the intriguing physicochemical properties of GDY, it also shows promise in various applications, such as water splitting cell^[7] and solar steam generation^[8].




Fig. 1. The synthesis and applications of graphdiyne.

References

- [1] J. Zhang *et al.*, *J. Am. Chem. Soc.* **137**, 24 (2015).
- [2] J. Zhang *et al.*, *Adv. Mater.* **28**, 1 (2016).
- [3] J. Zhang *et al.*, *Adv. Mater.* **29**, 19 (2017).
- [4] J. Zhang *et al.*, Submitted, 2018
- [5] J. Zhang *et al.*, *J. Phys. Chem. C* **120**, 19 (2016).
- [6] J. Zhang *et al.*, *Adv. Mater.* **29**, 18 (2017).
- [7] J. Zhang *et al.*, *J. Am. Chem. Soc.* **138**, 12 (2016).
- [8] J. Zhang *et al.*, *Chem. Mater.* **29**, 14 (2017).